© Society of Education, India ISSN 0976-4828 http://www.soeagra.com



REVIEW ARTICLE

Medicinal Plants for Diabetes Mellitus: A Traditional Approach

Jinesh Kumar and Manish Kumar

Department of Zoology, Agra College, Agra

ABSTRACT

Plants have been used in treatment of diabetes mellitus all over the world for centuries. Wide variety of plant derived active principles representing numerous classes of chemical compounds have shown potential for the use in treatment of diabetes. The present paper reviews medicinal plants with their mechanism of action and their pharmacological test results. Many studies have confirmed the benefits of medicinal plants with hypoglycemic effects in the management of diabetes mellitus. The effects of these plants may delay the development of diabetic complications and correct the metabolic abnormalities.

Keywords: Medicinal Plants, DM, Hypoglycemia, Hyperglycemia

INTRODUCTION

Plants have been the major source of drugs in Indian system of medicine and other ancient systems in the world. Charaka Samhita and Sushruta Samhita give extensive description on various medicinal herbs (1). Information on medicinal plants in India has been systematically organized (1-4). The World Health Organization expert committee on diabetes has listed as one of its recommendations that traditional methods of treatment of diabetes should be further investigated (5).

Diabetes mellitus (DM) is a metabolic disorder resulting from a defect in insulin secretion, insulin action, or both.1-4 Insulin deficiency in turn leads to chronic hyperglycaemia with disturbances of carbohydrate, fat and protein metabolism [6-10]. As the disease progresses tissue or vascular damage ensues leading to severe diabetic complications such as retinopathy, neuropathy, nephropathy, cardiovascular complications and ulceration.Thus, diabetes covers a wide range of heterogeneous diseases. Diabetes is the most common endocrine disorder and by the year 2010, it is estimated that more than 200 million people worldwide will have DM and 300 million will subsequently have the disease by 2025[11-18].

Among the challenging health conditions that are increasingly affecting human beings is diabetes. This is a disease condition caused by a metabolic disorder of the body systems as a result of chronic high blood sugar (hyperglycaemia). This sort of health challenge is often associated with disturbances in the metabolism of fat, carbohydrate and protein thereby causing defects in insulin activities and secretion due to the inability of the pancreas to produce enough insulin. This can also be as a result of the failure of the body cells to respond to already-produced insulin [19-25].

This health condition has led to several mortalities and morbidities with highest cases recorded in Asia and Africa. According to World Health Organisation, several actions can be put in place to lower the impact of diabetes on human health. These actions can be through adopting healthy lifestyles such as eating healthy balanced diets, engaging in physical exercises and activities, reducing the marketing of unhealthy food products and putting in place health systems that provide standard care and services for the affected individuals [24].

As diabetes continues to rise, the need to create awareness and adopt plans on how to reduce someone's risk of being affected, recognise the onset of the disease and determine the best mode of treatment are becoming highly important. There are several means of managing and treating diabetes, however, researchers reveal that natural remedies are more viable unlike the synthetic drugs and oral medications that may pose undesirable side effects to the body [20-24].

Even though herbal medicines produced from plants can be used as an anti-diabetic remedy alone, yet they can also be used alongside with other synthetic drugs such as insulin. The truth is that plant extracts from medicinal plants had been used over several years for treating numerous health disorders and diabetes is not an exception. Some key active components present in certain medicinal plants play significant roles in treating diseases and such powerful components continue to show promising effects in treating diabetes mellitus and any complications associated with it. We described below are some powerful Medicinal plants for treating diabetes.

COMMON ANTI-DIABETIC MEDICINAL PLANTS



Okra

Okra (Abelmoschus esculentus) [6]

Okra is a plant crop that contains anti-hyperglycemic properties. Several studies have supported the efficacy of fresh okra, soaked okra juice and powdered okra in tackling diabetes. Ben-Chioma et al., (2013) investigated the anti-hyperglycaemic activity of okra (Abelmoschus esculentus) fruits in alloxan-induced diabetic Wistarrats. The researchers grouped the animals into five; A, B, C, D, and E groups and they induced Diabetic Mellitus in groups B to E by single intravenous injection of alloxan, 65mg/kg body weight. However, group A was used as the control group (non-diabetic). Group C and group D diabetic rats were administered with dried powdered form and aqueous extracts of okra fruit respectively while group E was given a standard anti-diabetic drug (glibenclamide). The trial period was for 14 days after which, blood samples were taken and the fasting blood glucose (FBS) levels measured at days 3, 7, and 14 respectively. The study showed that both the dried powdered and aqueous form of okra significantly reduced the blood glucose levels of the animals at p<0.05. Therefore their results support the therapeutic use of okra as an anti-diabetic plant for managing and treating diabetes mellitus.



Allium cepa

Allium cepa [3, 4, 6, 21]

Allium cepa, which is popularly known as an onion is a spice plant that belongs to the Liliaceae family. From time immemorial, onion has been used for treating various types of diseases of which diabetes is part of them. Allium cepa helps to regulate the hypoglycemic activity which is associated with diabetes mellitus.

This is as a result of the present of flavonoids (quercetin) and sulphur compounds (S-methylcysteine) in Allium cepa which helps to reduce the level of blood glucose, lipid peroxidation, serum lipids as well as oxidative stress. These compounds also aid in insulin secretion as well as boost the antioxidant enzyme activities taking place in the body.

Onion extracts also aid in hypolipidemic activities. The hypolipidemic and hypoglycemic effects of onion is due to its ability to normalise the activities of HMG coenzyme-A reductase, liver hexokinase and glucose 6-phosphatase. Some preliminary clinical trials carried out reveal that glucose levels can be reduced by taking Allium cepa aqueous extracts. The hypoglycaemic and hypolipidemic abilities of Allium cepa are highly protective against the onset of hyperglycaemia, hyperlipidaemia and atherosclerosis, which are all associated with diabetes mellitus.



Milk Thistle

Milk Thistle [6]

Milk thistle is originally from the Mediterranean before spreading to other parts of the world. When the leaves are squeezed, they produce milky sap which suggests its name. Milk thistle belongs to the daisy family and it contains a polyphenol antioxidant compound known as silymarin that helps to control the blood sugar level.

Scientific studies reveal that milk thistle can be used for treating Type 2 diabetes. A scientific study carried out showed that diabetic patients who took silymarin for up to 4 months showed significant improvement in their glycemic profile. This also includes a significant decrease in fasting blood glucose, LDL cholesterol, glycosylated hemoglobin and triglycerides when compared to individuals taking only placebo.



Guggul [3, 4,]

Guggul is another name for Oleogum resin or Commiphora mukuland this is popularly used in Asia for producing herbal medicine for treating diabetes. Oleogum resin is a yellowish substance that is characterised by balsamic odour and each tree produces up to 900 g of resin. Guggul tree is originally from Bangladesh, Pakistan and India where it is mainly used in folk medicines. Guggulsterones are the active ingredient present in guggul.

Guggul extracts are highly effective in controlling the blood sugar level due to its role in glucose and lipid metabolism. Besides, guggul has a protective effect on pancreas which assists in the natural production of insulin in the body.

Commipheric acid and guggulipid present in guggul help to activate the peroxisome proliferator-activated receptor gamma (PPAR γ) hormone that regulates the overall development and functioning of fat cells which are vital for the healthy metabolism of cholesterol and glucose.



Loquat[6]

Loquat is botanically known as Eriobotrya japonica and belongs to the rose family. The loquat tree grows up to thirty feet with evergreen leaves that are between three to four inches wide and five to twelve inches long. Researchers reveal that the loquat leaf comprises of beneficial properties that can significantly influence the blood sugar level thus drinking loquat tea has been recommended as an ideal way of balancing the blood sugar level.

Loquat leaf contains polysaccharides and triterpenes chemicals especially tormentic acid that has been proven effective for boosting insulin production which significantly helps to lower the blood sugar.



Banaba Leaves

Banaba [6]

Banaba which is botanically known as Lagerstroemia speciosa (L.), belongs to the family of Lythraceae. Banaba is also known as queen's crape myrtle, the pride of India or queen's flower. Banaba is more common in the Philippines, India and Southeast Asia, where it is often used as an herbal medicine for treating diabetic patients.

Aqueous banaba leaf extracts can be used for lowering the blood sugar level and it's blood sugar lowering effect is sort of synonymous to that of insulin. Banaba leaf extracts help to transport glucose from the blood to the body cells.

But despite the beneficial impacts of banaba on blood sugar, diabetic patients are advised to be cautious when taking herbal medicines or supplements made with banaba.



Cinnamon Sticks

Cinnamon [3, 4, 6]

Cinnamon is a spice from the inner bark of many trees of the genus Cinnamomum which is widely used in food preparations. It has a characteristic golden-yellow colour with a hot aromatic taste, however, the pungent taste and scent are as a result of the presence of a cinnamic aldehyde or cinnamaldehyde and reaction with oxygen. It is a popular worldwide spice and many studies have been carried out to unravel the impact of cinnamon in reducing the blood glucose of diabetic patients.

Cinnamon does not contribute to caloric intake thus individuals with elevated glucose level or those with type 2 diabetes can benefit from the regular consumption of cinnamon. Moreover, those with high triglyceride or cholesterol level can benefit tremendously from the daily consumption of cinnamon as it aids in minimising inflammation and infection. Howbeit, some other researchers have contradictory claims that suggest that cinnamon is not truly effective in treating diabetes.



Garcinia Kola

Garcinia Kola [6]

Garcinia kola is a flowering plant that belongs to the Guttiferae or Clusiaceae family. Garcinia kola which

is also known as adi, Heckel, namijin, agbilu, orogbo, gworo, aki ilu, or bitter kola is anangiospermae that belongs to the family of Guttiferae.

Garcinia extract is highly nutritional and medicinal which suggests its wide usage in folk medicines for herbal medicines production. Garcinia possesses hypoglycaemic properties thus its aqueous extracts can be used for lowering blood sugar level. However, its hypoglycaemic effect is dependent on the dosage taken.

The hypoglycaemic properties of Garcinia are due to the present of flavonoid and other phytochemical compounds such as tannins, saponin and glycosides in Garcinia kola.



Garlic Cloves

Garlic [30-32]

Many researchers agree that garlic has hypoglycemic effects on bloodglucose, however, this claim is still disputable. Some experimental studies reveal that garlic and metformin treatment in diabetic patients for a period of 12 weeks lowered fasting blood glucose. Regular consumption of garlic extracts significantly reduces the blood glucose level.

The hypoglycemic effects of garlic on diabetes are as a result of the presence of volatile sulphur compounds, such as allyl mercaptan, alliin, allicin, ajoene, diallyl disulfide, diallyl trisulfide, S-allyl cysteine and diallyl sulfide in garlic. Garlic aqueous extracts are very effective for lowering insulin resistance in diabetic patients.



Green Tea [6]

The green tea leaves are very rich in antioxidants unlike other types of tea due to the way they are produced. Green teas are produced from the Camellia sinensis plant by steaming the fresh leaves from the plant.

Green tea is a common Asian beverage which has been reported to reduce or prevent the risk of type 2 diabetes mellitus as well as reduces insulin resistance. Green tea contains a high amount of polyphenols and Epigallocatechin-3-gallate that have strong anti-inflammatory and antioxidant effects on oxidative stress and inflammation associated with diabetes.



Licorice [3, 4, 6]

Licorice is the root of Glycyrrhiza glabra as well as a herbaceous perennial legume. The licorice root is an ancient traditional herbal remedy for tackling several ailments. Scientists reveal that licorice root from

the papilionaceae family may be effective for treating type 2 diabetes which is more common with overweight or obese individuals thus leading to insulin resistance.

Licorice roots contain anti-diabetic substances known as amorfrutins, which are capable of reducing the blood sugar levels as well as preventing inflammation associated with diabetics. The name amorfrutins was derived from the Amorpha fruticosa, which is a flowering plant belonging to the leguminous family.



Gymhema Sylvestre Leaves

Gymhema Sylvestre [3, 4, 6, 34]

Gymnema sylvestre is a tropical tree that belongs to the family of Apocynaceae and genus Gymnema. This tree is mostly prevalent in Africa, India, Australia, China and Vietnam. The leaf extract of Gymnemasylvestre is a powerful herbal medicine for tackling diabetes.

This is due to the presence of chemical components such as dihydroxy gymnemic triacetate, gymnemic acids I-VII, conduritol A and triterpenoid saponins (gymnemosides A-F and gymnemoside W1-2) in this plant. The effect of the Gymnema sylvestre leaf extracts is synonymous to 4 unit/kg of insulin thus highly beneficial for individuals suffering from diabetes mellitus.



Bitter Gourd

Momordica charantia[21-24, 32-37, 6]

Momordica charantia which is also known as karela, bitter gourd, balsam pear or bitter melon is a tendrilbearing vine used mainly for treating diabetes especially in India, the Caribbean, South America and East Africa.

Momordica charantia belongs to the family of Cucurbitaceae. Both the leaf extracts, fruit and seeds of bitter melon have hypoglycemic effects on diabetic patients. The key components of bitter melon which are responsible for its hypoglycemic effects are momordicin, stearic acid, charantin, eleostearic acid, insulin-like peptide [plant-(p)-insulin], oleanolic acids and cucurbutanoids. It is worthy to note that individuals who are allergic to the Cucurbitaceae family such as melons and gourds should apply caution when taking bitter melon fruit, seeds or leaf extracts.



Juniper berry [37, 38]

Juniper berries contain natural insulin and due to its ability to release insulin from the pancreas, researchers claim that it can alleviate hunger thus serves as a remedy for diet-controlled diabetes. Juniper

berry was reported to have antidiabetic and antihyperlipidemic effects on streptozotocin (STZ) nicotinamide induced diabetic rats. Moreover, the methanolic extract of this fruit showed a remarkable reduction in blood glucose levels in diabetic rats.

Having undergone several experimental studies, researchers agree that this fruit has hypoglycemic effects on diabetic patients.



Pterocarpus Marsupium

Pterocarpus Marsupium [30-38]

Pterocarpus marsupium is a large deciduous tree that belongs to the family of fabaceae (Leguminoceae). Parts of the Pterocarpus marsupium plant such as flowers, heartwood and leaves are used in Ayurveda due to the high medicinal properties present in this plant.

Pterocarpus marsupium which is also known as Bijasar, honne, kempu honne, Malabar Kino or Indian Kino is among the many powerful medicinal plants used for treating diabetes. The wood, bark and leaf extracts of P. marsupium possess anti-diabetic effects thus are used in folk medicine for treating diabetic patients.



Valeriana Wallichii

Valeriana Wallichii [35, 6]

Valeriana wallichii is a perennial herb that measures from 14 to 45cm height. This plant is a rhizome herb, a vital substitute for the European V. officinalis as well as belongs to the Valerianaceae family. It thrives very well mainly in the mountainous areas of the Himalayas.

The word Valeriana originated from Valere which connotes "to be in good health". This herb is very useful in Ayurvedic medicine as an antispasmodic, carminative, sedative, analeptic, nervine and stimulant. The Valeriana wallichii plant also possesses anti-diabetic properties thus can be used for treating individuals suffering from diabetes.



Yarrow Flower

Yarrow flower [21, 34]

Yarrow flower which is botanically known as Achillea millefolium is a flower with white petals that usually produces distinctive and pleasant odour. It is originally from Asia and Europe before spreading to other parts of the world. Yarrow flower contains inulin that produces fructose as a source of energy rather than glucose.

It is essential for diabetic patients because it doesn't cause a spike in blood glucose level after consumption. Due to its fibre effect, inulin helps to regulate bowel movement as well as aids in the elimination of waste products.



,

Cayenne Pepper [6]

Over the past years, cayenne pepper has been used traditionally for treating diabetes. A past study discovered that mice injected with capsaicin were healed of type 1 diabetes.

Type 1 diabetes, which is a more serious diabetic condition that starts in childhood due to the body's immune system attacking itself can be tackled with cayenne pepper. The study further revealed that injected capsaicin destroyed the pancreatic pain nerves thereby causing the body to start producing insulin normally. But being that cayenne is not a scientifically proven treatment of diabetes, no recommended dosage for consumption exists.

Because diabetes is a critical health condition, it is noteworthy for diabetic patients to continue taking cayenne alongside their prescribed medication in other to avoid complications. Some people are allergic to cayenne thus such individuals should desist from taking it. Excessive consumption of cayenne pepper can cause stomach irritation in some people thus should be consumed with care. Cayenne supplements can interact with certain types of medications such as blood thinners and aspirin [.



Fenugreek [2,5]

Fenugreek is botanically known as *Trigonella foenumgraecum* and it is an annual plant that belongs to the family of Fabaceae, whose seeds are widely used as a spice for culinary purposes. Fenugreek is multipurposeful as it can be used both as a spice (seeds), vegetable (microgreens, fresh leaves, sprouts), herb (dry/fresh leaves) or even for medicinal purposes.

This spice is highly medicinal and contains hypoglycemic properties thus can be used for preparing herbal medicines for treating diabetes.

Terminalia chebula Retz. (Combretaceae)

It has been widely used in diabetes in Ayurveda and is widely distributed in India. An herbal formulation containing T. chebula named TRIPHALA is traditional medicine for the treatment of diabetes. Antidiabetic and renoprotective effects of the chloroform extract of T. chebula Retz seeds in streptozotocin-induced diabetic rats was proved. It has potent renoprotective action [28].

Terminalia catappa Linn. (Combretaceae),

It is found throughout the warmer parts of India and called an Indian almond. The antidiabetic potential of petroleum ether, methanol and aqueous extract of T. catappa fruits on fasting blood sugar levels and serum biochemical analysis in alloxan-induced diabetic rats was performed. All the three extracts produced a significant antidiabetic activity at dose levels of 1/5 of their lethal doses. The extract may act by β -cells regeneration. The effect may be due to β -carotine in reducing diabetic complications like glycosylation in alloxan induced diabetic rats [29].

In general, there is very little biological knowledge on the specific modes of action in the treatment of diabetes, but most of the plants have been found to contain substances like glycosides, alkaloids, terpenoids, flavonoids etc. that are frequently implicated as having antidiabetic effects. The research for

alternate remedies (from the plant kingdom) for diabetes mellitus will continue all over the world as the disease poses many challenges not only to the physician but also to the researcher.

REFERENCES

- 1. WHO Expert committee on diabetes mellitus: second report, World Health Organ. Tech. Rep. Ser., 646, pp.1-80 (1980).
- 2. Diabetes Mellitus: Report of a WHO Study Group, World Health O Adesuyi AO, Elumm IK, Adaramola FB, Nwokocha AGM. (2012), Nutritional and Phytochemical Screening of Garcinia kola. Journal of Food Science and Technology 4(1), 9–14.
- 3. Nadkarni K.M.: Indian Materia Medica, 3rd ed., pp. 202-207 Popular Book Depot, Mumbai 1954. 13.
- 4. Kirtikar K.R., Basu B.D.: Indian Medicinal Plants, 2nd ed., pp. 1052-1054, Lalit Mohan Basu Publications, Allahabad 1933. 14.
- 5. Chopra R.W., Chopra I.C., Handa K.L., et al.: Indigenous Drugs of India, 2nd ed., pp. 314-316,
- 6. Dhar Kumar PJ, Clark M. Textbook of Clinical Medicine. Pub: Saunders (London), pp 1099-1121, 2002.
- 7. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the expert committee on the diagnosis and classification of Diabetes Mellitus. Diabetes Care 1997; 20: 1183-1197.
- 8. Beverley B, Eschwège E. The diagnosis and classification of diabetes and impaired glucose tolerance. In: Textbook of Diabetes 1 Ed: John C Pickup and Gareth Williams Third edition; Chapter 2, pp 2.1-2.11, 2003.
- 9. Lindberg G, Lindblad U, Melander A. Sulfonylureas for treating type 2 diabetes mellitus. Cochrane Database Systemic Reviews volume 3, 2004.
- 10. Bearse MA Jr, Han T, Schneck ME, et al. Local multifocal oscillatory potential abnormalities in diabetes and early diabetic retinopathy. Invest Ophthal Vis Sci 2004; 45 : 3259-3265.
- 11. Hove MN, Kristensen JK, Lauritzen T, Bek T. The prevalence of retinopathy in an unselected population of type 2 diabetes patients from Arhus County, Denmark. Acta Ophthalmol Scand 2004; 82: 443-448.
- 12. Seki M, Tanaka T, Nawa H, et al. Involvement of brain-derived neurotrophic factor in early retinal neuropathy of streptozotocin-induced diabetes in rats: therapeutic potential of brain-derived neurotrophic factors for dopaminergic amacrine cells. Diabetes 2004; 53: 2412-2419.
- 13. Moran A, Palmas W, Field L, et al. Cardiovascular autonomic neuropathy is associated with microalbuminuria in older patients with type 2 diabetes. Diabetes care 2004; 27: 972-977.
- 14. Huang C, Kim Y, Caramori ML, et al. Cellular basis of diabetic nephropathy: II. The transforming growth factorbeta system and diabetic nephropathy lesions in type 1 diabetes. Diabetes 2002; 51: 3577-3581.
- 15. Shukla N, Angelini GD, Jeremy JY, et al. Homocysteine as a risk factor for nephropathy and retinopathy in type 2 diabetes. Diabetologia 2003; 46; 766-772.
- 16. Svensson M, Eriksson JW, Dahlquist G. Early glycemic control, age at onset, and development of microvascular Bastaki S 128 complications in childhood-onset type 1 diabetes: a population-based study in northern Sweden. Diabetes Care 2004; 27: 955-962.
- 17. Saely CH, Aczel S, Marte T, et al. Cardiovascular complications in type 2 diabetes mellitus depend on the coronary angiographic state rather than on the diabetes state. Diabetologia 2004; 47: 145-146.
- 18. Wallace C, Reiber GE, LeMaster J, et al. Incidence of falls, risk factors for falls, and fall-related factures in individuals with diabetes and a prior foot ulcer. Diabetes Care 2002; 25: 1983-1986.
- 19. Centers for Disease Control and Prevention (CDCP). History of foot ulcer among persons with diabetes---- Unites States, 2000-2002. Morbidity & Mortality Weekly Report. 2003; 52: 1098-1102 and Sons Ltd., Calcutta 1958.
- 20. Ahmed I, Cummings E., Sharma A. K, Adeghate E and Singh J. (2004), Beneficial effects and mechanism of action of Momordica Charantia fruit juice in the treatment of streptozotocin-induced diabetes mellitus in rats. Mol Cell Biochem; 261:63-70.
- 21. Akash MS, Rehman K, Chen S. (2014), Spice plant Allium cepa: dietary supplement for treatment of type 2 diabetes mellitus, Nutrition journal 30(10):1128-37.
- 22. Ben-Chioma, A. E., Tamuno-Emine, D. G. and Dan, D. B. (2013), The Effect of Abelmoschus esculentus in Alloxan Induced Diabetic Wistar Rat, International Journal of Science and Research (IJSR), pp. 540-543.
- 23. Dans A. M, Villarruz M. V, Jimeno C. A, Javelosa M. A, Chua J and Bautista R. (2007), The effect of Momordica Charantia capsule preparation on glycemic control in type2 diabetes mellitus needs further studies. J Clin Epidemiol; 60:554-9.
- 24. Drucker D. J and Nauck M. A. (2006), The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. The Lancet;368(9548):1696–1705.
- 25. Follmann D, Elliott P, Suh I, (1992), Cutler J. Variance imputation for overviews of clinical trials with continuous response. J Clin Epidemiol; 45: 769–773.
- 26. Elisângela Düsman, Igor Vivian de Almeida, Ana Carolina Coelho, Thiago José Balbi, Lilian TatianiDüsman Tonin, and Veronica Elisa Pimenta Vicentini, "Antimutagenic Effect of Medicinal Plants Achillea millefolium and Bauhinia forficata In Vivo," Evidence-Based Complementary and Alternative Medicine, vol. 2013, Article ID 893050, 6 pages, 2013. doi:10.1155/2013/893050
- 27. Bahle S., John A.O.: Med. J. Islam. Acad. Sci. 13, 75 (2000).
- 28. Narayan N.S., Sastry K.N.V.: Mysore J. Agric. Sci. 9, 132 (1975).
- 29. Khan B.A., Abraham A., Leelamma S.: Ind. J. Biochem. Biophys. 32, 106 (1995).
- 30. Yadav S., Vats V., Dhunnoo Y., et al.: J. Ethnopharmacol. 82, 111 (2002).

- 31. Kesari A.N., Gupta R.K., Watal G.: J. Ethnopharmacol. 97, 247 (2005).
- 32. John L.S., John T. A., Lawrence A. L., et al.: J. Am. Coll. Nutr. 22, 524 (2003).
- 33. Bramachari H.D., Augusti K.T.: J. Pharm. Pharmacol. 13, 381 (1961).
- 34. Joao B.C., Santos A.R.S., Filho V.CJ, Yunes. R.A., 1998. Areview of the plants of the genus Phyllanthus: Their chemistry, pharmacology, and therapeutic potential. Medicinal Res. Rev. 18, 225-258.
- 35. Johnston A., 1987. Plants and the Blackfoot. Occasional Paper No. 15. Lethbridge Historical Society, Historical Society of Alberta, Alberta, Canada, 55 pp. 36. Kumar G.P., Sudheesh S., Vijayalakshmi N.R., 1993. Hypoglycemic effect of Coccinia indica: mechanism of action.
- Planta Medica 59, 330-332.
- 37. Kumari K., Augusti K.T., 2002. Antidiabetic and antioxidant effects of S-methyl cysteine sulfoxide isolated from onions (Allium cepa Linn) as compared to standard drugs in alloxan diabetic rats. Ind. J. Exp. Biol. 40, 1005-1009.
- 38. Maghrani M., Lemhadri A., Jouad H., Michel J., Eddouks M., 2003. Effect of the desert plant Retama raetam on glycaemia in normal and streptolotocin-induced diabetic rats. J. Ethnopharmacol. 87, 21-25.